

# Hour-1 Bundle

## Initial Resuscitation for Sepsis and Septic Shock



Initiate bundle upon recognition of sepsis/septic shock.

*May not complete all bundle elements within one hour of recognition.*

1

Measure lactate level.  
Remeasure lactate if initial lactate elevated ( $> 2$  mmol/L).

2

Obtain blood cultures before administering antibiotics.

3



Administer broad-spectrum antibiotics.

4

Begin rapid administration of 30 mL/kg crystalloid for hypotension or lactate  $\geq 4$  mmol/L.

5

Apply vasopressors if hypotensive during or after fluid resuscitation to maintain a mean arterial pressure  $\geq 65$  mm Hg.



Bundle: [SurvivingSepsis.org/Bundle](https://SurvivingSepsis.org/Bundle)

Complete Guidelines: [SurvivingSepsis.org/Guidelines](https://SurvivingSepsis.org/Guidelines)

## **HOURLY-1 BUNDLE: INITIAL RESUSCITATION FOR SEPSIS AND SEPTIC SHOCK:**

- 1) Measure lactate level.\*
- 2) Obtain blood cultures before administering antibiotics.
- 3) Administer broad-spectrum antibiotics.
- 4) Begin rapid administration of 30mL/kg crystalloid for hypotension or lactate  $\geq 4$  mmol/L.
- 5) Apply vasopressors if hypotensive during or after fluid resuscitation to maintain a mean arterial pressure  $\geq 65$  mm Hg.

\*Remeasure lactate if initial lactate elevated ( $> 2$  mmol/L).

# Surviving Sepsis Campaign®

1. \*Act quickly upon sepsis & septic shock recognition
2. Minimize time to treatment - sepsis & septic shock are medical emergencies
3. Monitor closely for response to interventions
4. Communicate sepsis status in hand-offs

\*All elements of the Hour-1 bundle may or may not be completed in the first hour after sepsis recognition

[survivingsepsis.org](http://survivingsepsis.org)

## HOURLY-1 BUNDLE: INITIAL RESUSCITATION FOR SEPSIS AND SEPTIC SHOCK:

- 1) Measure lactate level.\*
- 2) Obtain blood cultures before administering antibiotics.
- 3) Administer broad-spectrum antibiotics.
- 4) Begin rapid administration of 30mL/kg crystalloid for hypotension or lactate  $\geq 4$  mmol/L.
- 5) Apply vasopressors if hypotensive during or after fluid resuscitation to maintain a mean arterial pressure  $\geq 65$  mm Hg.

\*Remeasure lactate if initial lactate elevated ( $> 2$  mmol/L).



# Surviving Sepsis Campaign®

1. \*Act quickly upon sepsis & septic shock recognition
2. Minimize time to treatment - sepsis & septic shock are medical emergencies
3. Monitor closely for response to interventions
4. Communicate sepsis status in hand-offs

\*All elements of the Hour-1 bundle may or may not be completed in the first hour after sepsis recognition

[survivingsepsis.org](http://survivingsepsis.org)